ISSN (print) 0093-4666 © 2011. Mycotaxon, Ltd. ISSN (online) 2154-8889

MYCOTAXON

Volume 116, pp. 49–52 *April–June 2011*

DOI: 10.5248/116.49

Lecanora, Phaeophyscia and *Rinodina* species new to Turkey

Belgin Arslan, Şule Öztürk & Seyhan Oran*

Department of Biology, Faculty of Arts & Sciences, Uludag University, 16059, Bursa, Turkey
Correspondence to *: *seyhana@uludag.edu.tr*

Abstract — Four lichenized fungi (*Lecanora jamesii, L. juniperina, Phaeophyscia hirsuta,* and *Rinodina orculata*) are reported for the first time from Turkey. Comments on their habitat, substrate, and morphological and anatomical features are provided.

Key words — *Ascomycota*, Bursa, Çanakkale, lichens

Introduction

Lichenological studies in Turkey have rapidly increased in recent years. Prior to 2004, 361 papers referring to lichens were published (John 2004). Since then, many floristic studies have added several new records to the lichen biodiversity of Turkey (e.g. Yazıcı & Aslan 2005, Güvenç et al. 2006, John & Türk 2006, Halıcı et al. 2007, Yavuz & Çobanoğlu 2007, Candan & Türk 2008, Halıcı & Aksoy 2009, Kocakaya et al. 2009, Kınalıoğlu 2009, 2010). In spite of the increase in the number of studies, additional research is needed in some provinces where lichen biodiversity is still not sufficiently understood. This paper contributes further knowledge of the lichenized fungi in Turkey.

Materials & methods

Vouchers are stored in BULU (Herbarium of Uludag University, Sciences and Arts Faculty, Bursa, Turkey) and their accession numbers are given in parenthesis with the locality information. The specimens were examined with stereomicroscope (Prior), and light microscope (Krüss). Anatomical sections were examined in water, 10% KOH, and Lugol's iodine solution. Ascospore measurements were carried out in water and the extreme values outside the main range are given in parenthesis.

Species recorded

Lecanora jamesii J.R. Laundon 1963 Fig. 1

Specimen examined—**TURKEY. Çanakkale:** Bayramiç; Kazdağı Mountain, Dalaksuyu place, beech forest, 39°46N, 26°58E, alt. 1300 m, on *Fagus orientalis* Lipsky (*Fagaceae*), 18 Aug. 2005, leg. S. Oran, det. S. Oran (BULU 13785).

Thallus to 2–3 cm in diam., forming continuous, smooth or granular patches, grey; prothallus when present black; soredia arising in ± circular soralia, to 1 mm diam, farinose, pale yellow (to whitish on storage). Apothecia absent. Thallus P–, K–, C–, soralia sometimes weakly K+ yellowish.

On smooth bark of deciduous trees, most frequently *Salix* in damp situations, rarely on wood or siliceous rock; locally abundant (Purvis et al. 1992). In our study area, *L. jamesii* was found in a forest on smooth bark of *Fagus orientalis*.

Lecanora jamesii has been reported from western Europe and Chile (Purvis et al. 1992), and Iran (Valadbeigi & Sipman 2010). Clauzade & Roux (1985) and Purvis et al. (1992) describe the species in detail.

Lecanora juniperina Śliwa 2004 Fig. 2

Specimen examined—**TURKEY. Bursa:** İnegöl; road of Hilmiye–Oylat, 1 km from Hilmiye, rocky area, 39°56N, 29°35E, alt. 685 m, on *Juniperus* sp. (*Cupressaceae*), 19 Sept. 2004, leg. B. Arslan, det. Ş. Öztürk (BULU 9959).

Thallus granulose to areolate-rimose, grayish. Apothecia 0.3–1.2 mm in diam.; disc pale, sometimes yellow, usually pruinose, smooth; margin usually prominent, crenulate to dentate, paler than the disc; epihymenium granular, granules not soluble in K and N. Ascospores 9.0–13.5 × 4.5–6.0(–7.5) µm, ellipsoid.

Lecanora juniperina grows on tree bark (*Juniperus* and *Quercus*) and wood at intermediate elevations (Nash et al. 2004). Similar to the literature, our specimen was found on bark of *Juniperus* sp. in rocky area and was noted for its small thallus with very crowded and heavy pruinose apothecia.

Nash et al. (2004) and Śliwa (2007), who also describe the species in detail, recorded *L. juniperina* from North America. Valadbeigi & Sipman (2010) also report the species from Iran.

Phaeophyscia hirsuta (Mereschk.) Essl. 1978 Fig. 3

Specimen examined—**TURKEY. Bursa:** İnegöl; road of Çayyaka–Lütfiye, 1 km from Çayyaka, oak woodland, 39°59N, 29°28E, alt. 489 m, on *Juglans* sp. (*Juglandaceae*), 06 Jun. 2004, leg. B. Arslan, det. Ş. Öztürk (BULU 9601).

Thallus foliose, up to 3(–4) cm in diam., often orbicular; lobes 0.5–1(–1.5) mm broad, upper surface gray to gray-brown or brown, with cortical hairs near the lobe ends, sorediate; soralia labriform; lower surface black; rhizines simple, black. Apothecia numerous, up to 2 mm in diam, sessile, margin usually with cortical hairs.

Phaeophyscia hirsuta grows mostly on bark but also commonly on rock (Nash et al. 2004). Our specimen was collected from a trunk of *Juglans* sp. in a protected forest area.

Clauzade & Roux (1985), Nash et al. (2004), and Liška et al. (2008) have reported *P. hirsuta* from North America, Europe, and Africa. The first two papers describe the species in detail.



Fig. 1. *Lecanora jamesii* (BULU 13785). Fig. 2. *Lecanora juniperina* (BULU 9959). Fig. 3. *Phaeophyscia hirsuta* (BULU 9601). Fig. 4. *Rinodina orculata* (BULU 9818).

Rinodina orculata Poelt & M. Steiner 1970 Fig. 4

Specimen examined—**TURKEY. Bursa:** İnegöl; road of Kurşunlu–Güzelyurt, 3 km from Kurşunlu, oak woodland, 40°02N, 29°42E, alt. 575 m, on *Quercus* sp., 20 Jun. 2004, leg. B. Arslan, det. B. Arslan (BULU 9818).

Thallus crustose, usually effuse and discontinuous, composed of small verrucae, whitish, yellowish-brown to reddish-brown. Apothecia lecanorine, adnate to sessile, 0.2–0.4(–0.6) mm in diam., thalline margin thick, persistent. Apothecial cortex up to 20 µm, I+ blue; disc brown, plane; epihymenium red-brown. Ascospores 12–15(–17) × 6–8 µm, *Physconia*-type.

Rinodina orculata occurs at high altitudes in coniferous forests, where it grows mainly on small coniferous twigs. Our lichen sample, which was found on *Quercus* bark in a broad-leaved forest, was diagnosed by its numerous and round apothecia with dark brown discs.

This is a montane species with a distribution centered in the European Alps where it mostly grows on *Rhododendron* spp. Giralt (2001), who describes the species in detail, has reported the species from the Iberian Pyrenees and Portugal; *R. orculata* has also been recorded in the Czech Republic (Liška et al. 2008).

Acknowledgements

We would like to thank Ayşen Türk (Eskişehir, Turkey) and M. Gökhan Halici (Kayseri, Turkey) for reviewing this paper and Dr. Harrie Sipman (Berlin) for checking the specimens and providing valuable suggestions for the manuscript. Collection of *Lecanora jamesii* was made possible by project (2006/63), funded by the Unit of Scientific Research Projects, Uludag University.

Literature cited

Candan M, Özdemir Türk A. 2008. Lichens of Malatya, Elazığ and Adıyaman provinces (Turkey). Mycotaxon 105: 19–22.

Clauzade G, Roux C. 1985. Likenoj de Okcidenta Eŭropo. Ilustrita Determinlibro. Royan. Bulletin de la Société Botanique du Centre-Ouest Nouvelle série–Numéro Spécial.

Giralt M. 2001. The lichen genera Rinodina and Rinodinella (lichenized Ascomycetes, Physciaceae) in the Iberian Peninsula. Bibliotheca Lichenologica, Band 79, Stuttgart.

Güvenç Ş, Öztürk Ş, Aydın S. 2006. Contributions to the lichen flora of Kastamonu and Sinop provinces in Turkey. Nova Hedwigia 83: 67–98. doi:10.1127/0029-5035/2006/0083-0067

Halıcı MG, Aksoy A. 2009. Lichenized and lichenicolous fungi of Aladağlar National Park (Niğde, Kayseri and Adana Provinces) in Turkey. Turkish Journal of Botany 33: 169–189. doi:10.3906/bot-0810-14

Halıcı MG, Candan M, Özdemir Türk A. 2007. New records of lichenicolous and lichenized fungi from Turkey. Mycotaxon 100: 255–260.

John V. 2004. Lichenological studies in Turkey and their relevance to environmental interpretation. Abstract book, XI OPTIMA meeting, 5.–11.9.2004 Belgrad: 45.

John V, Türk A. 2006. Species/area curves for lichens on gypsum in Turkey. Mycologia Balcanica 3: 55–60.

Kınalıoğlu K. 2009. Lichens from the Amasya, Çorum, and Tokat regions of Turkey. Mycotaxon 109: 181–184.

Kınalıoğlu K. 2010. Lichens of Ordu Province, Turkey. Mycotaxon 112: 357–360. doi:10.5248112.357

Kocakaya M, Halıcı MG, Aksoy A. 2009. Lichens and Lichenicolous Fungi of Kızıldağ (Derebucak, Konya). Turkish Journal of Botany 33: 105–112. doi:10.3906/bot-0810-2

Liška J, Palica Z, Slavíková Š. 2008. Checklist and Red List of lichens of the Czech Republic. Preslia 80: 151–182.

Nash III TH, Ryan BD, Diederich P, Gries C, Bungartz F. 2004. Lichen Flora of the Greater Sonoran Desert Region, Volume 2. Tempe, Arizona State University.

Purvis OW, Coppins BJ, Hawksworth DL, James PW, Moore DM. (Eds.) 1992. The lichen flora of Great Britain and Ireland. London, Natural History Museum Publications.

Śliwa L. 2007. A revision of the *Lecanora dispersa* complex in North America. Polish Botanical Journal 52(1): 1–70.

Valadbeigi T, Sipman HJM. 2010. New records of lichens and lichenicolous fungi from Iran and their biogeographical significance. Mycotaxon 113: 191–194. doi: 10.5248/113.191

Yavuz M. Çobanoğlu G. 2007. Lichen flora of Pamukkale (Hierapolis) Turkey. Pakistan Journal of Biological Sciences 10(17): 2998–3001.

Yazıcı K, Aslan A. 2005. Six new lichen records from Turkey. Mycotaxon 93: 359–363.